

Evidence for a Very Broad Line Region in PG 1138+222

*Paolo Marziani & Jack W. Sulentic*¹

Department of Physics & Astronomy
University of Alabama
Tuscaloosa, Al 35487-0324

U.S.A.

Received:

Accepted:

¹Visiting Astronomer, Kitt Peak National Observatory

Abstract

We present a high S/N spectrum for the quasar PG 1138+222. We detect a very broad HeII λ 4686 emission line component with twice the FWHM of the conventional broad line region (BLR) as evidenced by the Balmer lines. The profile shape and centroid redshift also distinguish this HeII component from the BLR features. The large ratio of HeII λ 4686 to any analogous H β emission component is an indicator that it arises in a very high density region ($n_e \gtrsim 10^{11} - 10^{13} \text{ cm}^{-3}$). This Helium component is probably emitted in a Very Broad Line Region (VBLR), where the radiation field is so strong that the Strömgren depth becomes similar to the geometrical thickness of the emitting clouds. The gas could therefore be optically thin to the Lyman continuum.

Subject headings: Galaxies: nuclei – Line Formation – Line Profiles – Quasars: Spectroscopy

1 Introduction

Photoionization models are fairly successful in reproducing the broad line spectra of Seyfert galaxies and quasars. Recent attempts to compute the emergent spectrum of the Broad Line Region (BLR) have led to significant improvements because a wider range of electron density has been considered (n_e from the “canonical” value 10^9 up to 10^{13} cm^{-3}) (Rees, Netzer & Ferland, 1989; Dumont & Collin-Souffrin, 1990; Ferland et al. 1992). It is still unclear whether density decreases smoothly with distance from the continuum source or whether the low ionization lines (Balmer lines, $\text{MgII}\lambda 2800$, FeII emission) and the high ionization lines ($\text{Ly}\alpha$, $\text{CIV } \lambda 1549$, and also $\text{HeII } \lambda 4686$) are emitted in two different environments (with different n_e and probably different underlying kinematics). The possibility that LIL and HIL arise in different environments could account for the apparent blue shift of the HIL lines relative to the LIL (Gaskell 1982). Such a difference, for example, arises naturally in a radiating accretion disk model. The accretion disk might be the LIL emitter, provided that it is illuminated by a significant amount of back-scattered continuum radiation (Dumont & Collin-Souffrin, 1990). One problem with this interpretation is the poor overlap between model predictions and the observed properties of the emission lines (Sulentic et al. 1990).

Irrespective of any multi-zoned structure or stratification it is likely that extreme physical conditions will be encountered at the inner edge of the BLR. The wings of the Balmer lines and some HIL could arise in a *Very Broad Line Region* (VBLR) whose properties differ markedly from those of a composite BLR. Although high S/N spectra of Seyfert galaxies often show very extended and faint wings with FWZI up to $30,000 \text{ km s}^{-1}$ (see e.g. Akn 120 & IC 4329A, Marziani, Calvani & Sulentic, 1992) the possibility of a VBLR has only been discussed for Mkn 590 (Ferland, Korista & Peterson, 1990) and NGC 5548 (Peterson & Ferland, 1986; Wamsteker et al. 1990) as far as we are aware. The former authors introduced the idea of an optically thin VBLR for Mkn 590 in order to explain the absence of variability in the wings of the Balmer lines. A tenfold increase in the strength of the $\text{HeII}\lambda 4686$ line in NGC 5548, along with a much smaller increase in $\text{H}\beta$, was similarly ascribed to optically thin gas very

close to the continuum source (Peterson & Ferland, 1986). In some objects (where the OI λ 8446 line is observed) the wings of the Balmer lines are found to be stronger or more extended than those for OI. This line is believed to arise from gas that is optically thick to the Lyman continuum (Morris & Ward, 1988). This raises the possibility that the wings of the Balmer lines arise in an optically thin region. Ferland, Korista & Peterson (1990) proposed that a “screen” of optically thin gas located between the canonical BLR and the central continuum source might even reprocess part of the nonthermal continuum and create the *Big Bump*. The standard explanation for the Big Bump involves optically thick gas, either in an accretion disk, or in dense, filamentary structures (Lightman & White, 1988).

Little attention has been paid to the HeII λ 4686 feature as a valuable indicator of the range of properties inside the BLR. This is probably because this feature is usually contaminated by FeII emission. We find a very broad HeII λ 4686 component in PG 1138+222. This feature could only be detected with : (1) high S/N data, (2) no significant contamination from an underlying galaxy and (3) weak FeII emission between H β and H γ . Thus PG 1138+222 provides a rare opportunity to make a reliable study of the HeII λ 4686 profile. Our analysis of the HeII λ 4686 and Balmer line profiles are reported in this paper along with a heuristic analysis of the physical condition for gas in the VBLR.

2 Observations and results

2.1 Observations and data reduction

PG 1138+222 ($\equiv$ WAS 26) was observed at Kitt Peak National Observatory on February 19, 1991, with the Gold Camera attached to the Cassegrain focus of the 2.1 m telescope. A dispersion of 1.3 Å/pixel was obtained with a 600 l/mm grating and an 800 $\times$ 800 pixel TI chip. The spectral range covered was $\approx$ 4500–5500 Å. The slit width was $\approx$ 2.5 arcsec (yielding a spectral resolution of $\approx$ 3.2 Å FWHM), while the seeing was estimated to be $\approx$ 1.5 arcsec. Standard IRAF procedures were adopted to convert the observed spectrum to linear wavelength scale. Conversion to absolute flux units was based upon observations of the star

Feige 34.

2.2 Results

Figure 1 shows the spectrum of PG 1138+222. The redshift, computed from all identified narrow emission lines (reported in Table 1) is $z = 0.0632 \pm 0.0004$. Fluxes of the lines, uncorrected for galactic reddening, (Burstein & Heiles (1984) give $E(B - V)_{gal} = 0.0$) are reported in the second column of Table 1. The underlying continuum was first subtracted with a third-order spline function in order to estimate the relative contribution of the broad and narrow components. The broad components of $H\gamma$, $HeII\lambda 4686$ and $H\beta$ were then fitted with appropriate spline functions. Results are shown plotted on the same wavelength scale in Figures 2a b and c respectively. Absolute fluxes of the broad and narrow components are estimated to be accurate within $\pm 20\%$.

The narrow line component ratio $H\gamma/H\beta \approx 0.49 \pm 0.06(2\sigma)$ agrees (within the uncertainties of the measurement) with the value appropriate for the Narrow Line Region (NLR) as computed by Ferland & Netzer (1983). Several Seyfert galaxies and quasars show a ratio that is considerably larger. This suggests that it is not always advisable to infer the reddening from this ratio alone. The lack of other indicators leads us to assume negligible internal reddening in PG 1138+222 as a first approximation.

An interesting result not directly related to the main topic of this paper is the remarkable strength of the well resolved $[OIII]\lambda 4363$ line. Accepted at a face value, the ratio $[OIII]\lambda\lambda 4959,5007/[OIII]\lambda 4363$ implies an electron temperature of $T_e \approx 3.5 \times 10^4$ °K for an electron density $n_e = 10^4$ cm $^{-3}$ or $T_e \approx 2.8 \times 10^4$ °K for $n_e = 10^5$ cm $^{-3}$ (Osterbrock, 1988). The ratio $[OIII]\lambda 4363/H\beta_{NC} \approx 0.44$ and the implied T_e are too large to be consistent with photoionization. This ratio is too high even for gas whose abundance is a factor of ten below the solar value and with very high ionization parameter ($\Gamma \sim 0.01 - 0.1$, Ferland & Netzer 1983). Heating mechanisms other than photoionization (e.g shocks) are probably at work in the NLR of PG 1138+222. A complication connected with the $[OIII]$ line ratio given above involves the observation that $\lambda 4363$ profile is nearly twice as broad as $\lambda 5007$. The greater

FWHM of [OIII] λ 4363 is clearly seen in Figure 2a and may indicate structure in the NLR. Gas with n_e between 10^6 and 10^7 cm $^{-3}$ may contribute to the [OIII] λ 4363 emission. A higher density gas would produce a higher velocity dispersion which might explain the unusual ratio without invoking shocks.

The principal result of this paper is the detection of a broad component underlying the narrow HeII λ 4686 emission feature. It is 2.4 times broader than the mean FWHM for H β and H γ . Similar values of FWZI are reported in Table 1 for all three lines. However the FWZI measures for the Balmer lines are affected by the presence of a weak red wing extending up to $v_r \sim 7000$ km s $^{-1}$, which is not seen in HeII.

The FeII emission in PG1138+222 is weak, but lines from multiplets 27, 42 and 36 are likely contributors to the extended red wings of H γ and H β : FeII λ 5169 (multiplet 42) is clearly visible at the red edge of our spectrum. Moreover, the red wing of H γ is so weak, that even FeII emission slightly above noise could reduce or cancel out this feature. Nevertheless, we believe that the red wing is real because: (1) it is seen in both the H β and H γ profiles of our spectrum, (2) the S/N ratio of our spectrum is very high (≈ 50) and (3) Osterbrock & Shaw (1988) published a low resolution spectrum of PG 1138+222 where the red wing is also visible in H α .

The Balmer line profiles can be classified as type AR,R (i.e., red asymmetric profile with redshifted peak) in the scheme proposed by Sulentic (1989). This type is rather common among quasars and Seyfert galaxies. The BLR component of H β shows a redshift (at FWHM) of 400 ± 75 km/s relative to the narrow line component. The profile of HeII λ 4686, appears to be markedly different from the Balmer lines. This is clearly seen in the Figure 2a,b comparison between H γ and HeII which have comparable fluxes. More quantitatively, if we take the ratio FWHM to FWZI as an indicator of the “boxiness” of a profile, we find 0.25 and 0.28 for H β_{BC} and H γ_{BC} while for HeII λ 4686, the ratio is 0.67. The peak and FWHM centroid values for the broad HeII component show the same redshift as the NLR with an uncertainty of 300 km s $^{-1}$.

Another comparison that emphasizes the difference between the HeII and H β profiles involves the ratio between HeII λ 4686_{BC} and H β _{BC} as shown in Figure 3. The ratio was computed after subtraction of the underlying continuum and NLR components as well as smoothing with a gaussian filter. The error bars displayed in Fig. 3 represent the uncertainty due to different, extreme choices of the underlying continuum. They should be regarded as representative of the maximum error. The ratio HeII λ 4686_{BC}/H β _{BC} is about 0.2–0.3 in the central part of the profile (similar to the flux ratio integrated over the entire line profile). The ratio increases up to a value greater than one at $v_r \approx -2000 \text{ km s}^{-1}$, and it remains ~ 2 up to $v_r \approx -4000 \text{ km s}^{-1}$, where both H β and HeII fall below the 3σ noise level. Although the ratio also tends to increase on the red side, its maximum value is lower, namely ≈ 0.8 , because of the red wing on H β . If we interpret the red wing as a distinct component or, at least, as arising in a region different from HeII λ 4686, the values on the blue side can be assumed to give a more relevant estimate for the ratio. The profile ratio involving the red side of H β will also increase after correction for FeII contamination.

2.3 A comparison of line ratios and the implications

The average ratio between the equivalent width of HeII λ 4686 and H β is 0.095 ± 0.10 for the 87 quasars studied by Boroson & Green (1992). These EW measures include both the broad and narrow line emission. The NLR contribution is usually small for H β while the situation for HeII λ 4686 is less clear. The ratio of the broad components in PG 1138+222 is $F(\text{HeII}\lambda 4686)/F(\text{H}\beta) \approx 0.33$. This value is rather large, and in fact only 10 quasars in the above sample show $F(\text{HeII}\lambda 4686)/F(\text{H}\beta) \approx 0.20$; seven of these show $F(\text{HeII}\lambda 4686)/F(\text{H}\beta) \approx 0.30$. Only PG 1244+026 (0.51) and PG1534+580 (0.40) show stronger HeII than PG 1138+222. PG1534 shows a very broad HeII profile similar to PG1138+222 with little or no contamination from FeII. The contamination from FeII emission in PG 1244+026 is too strong to make reliable inferences. The ratio for PG 1138+222 is actually more extreme than this comparison suggests. We are comparing the strengths of composite features that arise in different regions. It is likely therefore that the ratio of the broad HeII component to any

analogous one in $H\beta$ is ≥ 2.0 .

A ratio of $\text{HeII}\lambda 4686_{BC}/H\beta_{BC} \geq 2-3$ is larger than case B situations (≈ 0.8), and much larger than the value expected for the BLR (~ 0.2), where $H\beta$ is collisionally enhanced by a factor of ≈ 5 . It is a strong indicator of high density ($n_e \sim 10^{11} - 10^{13} \text{ cm}^{-3}$) inside the BLR. Earlier studies (Shuder, 1982; Osterbrock & Shuder, 1982) based on the $\text{HeI}\lambda 5876/H\beta$ and $H\alpha/H\beta$ profile ratios showed that the electron density and/or the ionization parameter increased toward higher radial velocities. Evidence for the VBLR rests with differences in the width and shape of the broad $\text{HeII}\lambda 4686$ component compared to the Balmer lines. The $\text{HeII}\lambda 4686$ and $\lambda 1640$ lines in NGC 5548 (Wamsteker et al. 1990) were also found to be substantially broader than $H\beta$. The greater width of the HeII component points toward a velocity regime nearer the central engine than for the conventional BLR.

2.4 The formation of HeII lines

The formation of the $\text{HeII}\lambda 4686$ line has been thoroughly discussed by MacAlpine (1981), Grandi (1983), and MacAlpine et al. (1985). MacAlpine discussed the use of the $\text{HeII}\lambda 1640/\text{HeII}\lambda 4686$ ratio as a reddening indicator for the BLR, and suggested that these lines are likely to have a ratio close to the recombination value. The $\text{HeII}\lambda 4686$ line originates from the level $4 \rightarrow 3$ transition as does $\text{Pa}\alpha$. Mechanisms which can enhance the $\text{HeII}\lambda 4686$ line over its recombination value are: (a) collisional excitation from level 2 to 4 and (b) radiative pumping of level 4 due to diffuse $\text{Ly}\alpha$ fluorescence, either from $\text{Ly}\alpha$ photons generated inside the cloud or coming from other emission line clouds. The effectiveness of these processes depends on the He^+ population of level 2 ($n_{\text{He}^+}(2)$), which depends almost entirely on the diffuse $\text{HeII}\text{Ly}\alpha$ (i.e. $\text{HeII}\lambda 304$ photons) field. A reliable estimate of $n_{\text{He}^+}(2)$ is difficult to compute, because of the complex transfer properties of the $\text{HeII}\text{Ly}\alpha$ line. It is well known that the $\text{HeII}\text{Ly}\alpha$ photons can be destroyed through the OIII Bowen fluorescence mechanism, and by ionization of neutral hydrogen atoms (e.g. MacAlpine 1981; Netzer, Elitzur & Ferland, 1983).

It is beyond the scope of the present work to attempt a quantitative prediction which

would require detailed photoionization calculations. Nevertheless, if we assume that a very high density gas ($n_e \sim 10^{11} - 10^{13} \text{ cm}^{-3}$) is contributing to the emission it is inescapable that the HeII λ 4686 line will be strongly enhanced over case B. MacAlpine (1981) computed the number of resonance scatterings by HeII Ly α needed to sustain a given value of $n_{He^+}(2)$ where the strength of HeII λ 4686 is due equally to HI Ly α pumping and recombination. At densities $n_e \sim 10^{11} - 10^{13} \text{ cm}^{-3}$, equations (5) and (10) of MacAlpine (1981) show that the number of scatterings permits a significant contribution from collisional excitation to level 4 and, to an even greater extent, from radiative pumping. The strength of HeII λ 4686 is enhanced by a large factor over its case B value if the pumping is due to HI Ly α emitted from other emission line clouds. At densities $> 10^{11} \text{ cm}^{-3}$, the strength of H β is also lowered by collisional deexcitation further contributing to an increase of the HeII λ 4686_{BC}/H β _{BC} ratio.

Generally speaking, the stronger the He⁺ ionizing radiation field, the stronger the diffuse HeII Ly α field should become, thereby increasing the population of level 2 in the He⁺ atom. Other relevant effects that enhance HeII λ 4686 over its recombination value depend strongly on density. The values $\Gamma \sim 0.1$ and ~ 0.01 for $n_e \sim 10^{11}$ and $\sim 10^{13} \text{ cm}^{-3}$ respectively, should be regarded as the lower limits necessary to explain our profile ratio. Larger values of the ionization parameter (i.e. the strength of the radiation field to which the clouds are exposed if the density is fixed) would create more favourable conditions for large values of the HeII λ 4686/H β ratio.

The detailed photoionization calculations by Rees, Netzer & Ferland (1989) indeed show that the ratio HeII λ 4686/H β becomes larger than one if $n_e \sim 10^{13} \text{ cm}^{-3}$ and if the ionization parameter $\Gamma = Q(H^0)/4\pi r^2 n_e$ is greater than 0.01, where $Q(H^0)$ is the number of ionizing photons. A ratio of around three can also be obtained for $n_e \sim 10^{11} \text{ cm}^{-3}$ and $\Gamma \sim 0.1$ (cf. Fig. 5 of Ferland et al. 1992). Generally speaking, a stronger radiation field gives a stronger diffuse HeII λ 304 field, and hence a larger population of He⁺ level 2. Values of Γ larger than those given above (i.e., a stronger radiation field for densities $\gtrsim 10^{11} \text{ cm}^{-3}$) would create even more favorable conditions to increase the HeII λ 4686/H β ratio. This problem deserves more

attention from the theoretical point of view. Nevertheless in light of the previous analysis and the photoionization results, it is probably appropriate to assume that the density at the innermost edge of the VBLR is in the range $\sim 10^{11} - 10^{13} \text{ cm}^{-3}$. In the following calculations, we will consider two cases: (1) $n_e \sim 10^{13} \text{ cm}^{-3}$ and $\Gamma \sim 0.01$, and (2) $n_e \sim 10^{11} \text{ cm}^{-3}$ and $\Gamma \sim 0.1$, as representative of the physical conditions in the VBLR.

2.5 Estimate of the VBLR properties for PG 1138+222

The distance of the VBLR from the central continuum source can be written as $R_{in} \approx 1.58 \times 10^{16} (\Gamma n_e)_{11}^{-1/2} \text{ cm}$. The number of ionizing photons $Q(\text{H}^0)$ needed to compute Γ was estimated by extrapolating the optical continuum up to $h\nu = 30 \text{ keV}$ assuming an ionizing continuum of the form given by Netzer (1990). We are unaware of any far UV or X-ray observations that might help us here.

The filling factor f_f of the total HeII emitting region was estimated writing the luminosity of the HeII λ 4686 line as:

$$L(\text{HeII}\lambda 4686) = \int n_e n_{\text{He}^{++}} \alpha_B^{eff}(\text{HeII}\lambda 4686) h\nu 4\pi r^2 dr f_f, \quad (1)$$

where $\alpha_B^{eff}(\text{HeII}\lambda 4686)$ is the effective recombination coefficient (case B) for HeII and the integral is to be evaluated from R_{in} . We used $H_0 = 100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ throughout the paper. The electron density was assumed to scale as $n_e = n_{e0} (r/R_{in})^{-2} \text{ cm}^{-3}$. We considered $n_{e0} = 10^{11}$ and 10^{13} cm^{-3} (as in model B of Rees, Netzer & Ferland, 1989). The recombination coefficient is $\alpha_B^{eff}(\text{HeII}\lambda 4686) \approx 1.43 \times 10^{-13} \text{ cm}^3 \text{ s}^{-1}$ at $T_e \approx 2 \times 10^4 \text{ }^\circ \text{ K}$ (Osterbrock, 1988). If emission is confined in a shell of depth $0.1 R_{in}$, and if f_f does not depend on r , then $f_f \sim 2.6 \times 10^{-6} n_{e0,12}^{-2} R_{in,16}^{-3}$.

The covering factor (in the case of spherical geometry) is $f_c = R_{in} n_e f_f / 3 N_c \approx 0.034 R_{in,16} n_{e,12} f_{f,-6} N_{c,23}$, where N_c is the column density. A broad upper limit to the the covering factor can be obtained from the relation

$$L(\text{HeII}\lambda 4686) = h\nu [\alpha_B^{eff}(\text{HeII}\lambda 4686) / \alpha_B^{eff}(\text{He}^{++} \rightarrow \text{He}^+)] f_c Q(\text{He}^+), \quad (2)$$

valid if the gas is optically thick for photons with energy higher than 54.4 eV. The value of f_c for the HeII emitting gas should thus be $f_c \lesssim 0.29Q_{54}^{-1}(He^+)$ ($n_e \sim 10^{13} \text{ cm}^{-3}$), and $0.39Q_{54}^{-1}(He^+)$ ($n_e \sim 10^{11} \text{ cm}^{-3}$), where the number of He^+ photons is in units of 10^{54} . Ferland, Korista & Peterson (1990) suggest that the optically thin gas in Mkn 590 is distributed in a screen that completely covers the central source. In their view, the central source of power-law radiation is surrounded by ionized and partially transparent gas. However employing a *bare* power-law $f_\nu \propto \nu^{-\alpha}$ with $\alpha = 1.0$ to compute $Q(H^0)$ we obtain an upper limit to $f_c \lesssim 0.5 - 0.6$ for PG 1138+222. This suggests that there is no need to have complete coverage of the continuum source of photons above the He^+ threshold in order to account for the observed HeII λ 4686 luminosity.

Strong variations in HeII λ 4686 coupled with a weaker variation in the Balmer lines suggested that the HeII emitting gas in NGC 5548 (Peterson & Ferland, 1986) was optically thin to the Lyman continuum. The same conclusion probably holds for PG 1138+222. The depth of the He^{++} Strömgen sphere is $R_{He^{++}} = 3\Gamma(He^+)c/n_{He^{++}}\alpha_B^{eff}(He^{++} \rightarrow He^+)$, where $\alpha_B^{eff}(He^{++} \rightarrow He^+)$ is the total effective recombination coefficient, and $\Gamma(He^+)$ is the ionization parameter for He^+ , that is $\Gamma(He^+) = Q(He^+)/4\pi n_e c R_{in}^2$ (which is $\sim 0.1 \Gamma(H^0)$ with our choice of the continuum). The number of He^+ photons was again estimated assuming that the UV – X-ray continuum had the form tabulated by Netzer (1990). Assuming $n_{He^{++}} \sim 0.1n_e$, and using the recombination coefficient computed by Hummer & Storey (1987), we find $R_{He^{++}} \sim 6.5 \times 10^{10} \text{ cm}$ ($n_e \sim 10^{11} \text{ cm}^{-3}$) and $R_{He^{++}} \sim 4 \times 10^7 \text{ cm}$ ($n_e \sim 10^{13} \text{ cm}^{-3}$).

The Strömgen radius for H^+ is analogously $R_{H^+} = 3\Gamma(H^0)c/n_e\alpha_B^{eff}(H^+ \rightarrow H^0)$, and corresponds to $\sim 6 \times 10^8$ and $\sim 6 \times 10^{11} \text{ cm}$ for $n_e = 10^{13}$ and 10^{11} cm^{-3} respectively. If $n_e \sim 10^{11} \text{ cm}^{-3}$, the canonical value for the column density of active nuclei, $N_c \sim 7 \times 10^{22} \text{ cm}^{-2}$ is obtained at $R \approx 7 \times 10^{11} \text{ cm}$. This suggests that the HeII emitting gas is most probably optically thin to the Lyman continuum. However, if $n_e \sim 10^{13} \text{ cm}^{-3}$, Γ would have to be ~ 0.1 in order for the Strömgen depth to become comparable to the geometrical

thickness of the clouds. Furthermore, the canonical value of N_c given above refers to neutral gas. It is probably a reliable estimate of the total cloud N_c under standard BLR conditions, where the Strömngren radius should be much smaller than the geometrical thickness of the clouds. The simplest choice was to also adopt this value of the column density for the VBLR.

2.6 Further considerations

Great care should be taken in using HeII λ 1640/HeII λ 4686 ratio as a reddening indicator. If the line profile of HeII λ 4686 is significantly broader than that of H β , as is the case for PG 1138+222 and as may be the case for most HeII-strong objects, emission in the line wings could give a non-negligible contribution to the total flux. As outlined above, there are reasons to believe that emission in the wings comes from high density gas in a VBLR exposed to an extremely strong radiation field. In these conditions, the HeII λ 4686 is expected to be strongly enhanced with respect to case B of recombination theory and to remain optically thin, while the optical depth in the HeII λ 1640 line could become larger than 1 (Ferland et al. 1992). Differences in the enhancement of HeII λ 4686 and λ 1640 may depend primarily on the relative contributions of clouds of different densities (exposed to different radiation field intensities), whose distribution inside the emitting region is unknown.

There is a kinematic difference between our VBLR evidence and that from Ferland et al. (1990). Our broad HeII λ 4686 component is centered at or near the systemic (as indicated by the NLR) velocity of the quasar. Their broad H β component shows a significant redshift $\Delta V \approx 10^3$ km/s. The difference in redshift between the HeII feature and the underlying Balmer features suggests that both cannot arise in the same kind of optically thin region. A similar comment applies to Wamsteker et al (1990) where the (VBLR) HeII λ 4686 feature in NGC 5548 shows a 2×10^3 km/s blueshift. We would only expect a gravitational redshift in the simplest static model for a VBLR. The differing velocity results suggest that obscuration and/or radial motion effects are present in the VBLR.

3 Conclusion

The introduction of a VBLR is not motivated simply for taxonomical convenience. The heuristic value of the VBLR stems from the physical conditions of the gas located closest to the central continuum source: as the distance decreases, the depth of the Strömgren radius also increases, making the gas optically thin to the HI Lyman continuum. We suggest that the large value of the ratio $\text{HeII}\lambda 4686_{BC}/H\beta_{BC} > 1$ can be explained if the VBLR is made up of dense clouds with $n_e \sim 10^{11} - 10^{13} \text{ cm}^{-3}$. If the above assumptions are true the HeII emission arises in region that may not reprocess most of the continuum emission. Thus, it could not account for the UV emission from the “big bump” (Ferland et al. 1990). This filtering environment might not be the “warm absorber” suggested by X-ray observations (Yaqoob, Warwick & Pounds 1989).

Other points remain to be clarified. The first is whether the quasars with strong HeII emission should be regarded as the high-end tail of a distribution or as a different population of objects. It is also unclear whether the large spread in the ratio $\text{HeII}\lambda 4686_{BC}/H\beta_{BC}$ ($\sim 0 - 0.88$ among the Seyfert 1 galaxies studied by Osterbrock, 1977) reflects inclination effects, the distribution of matter inside the BLR, or, less probably, intrinsic differences in the ionizing continuum. Since the VBLR is the innermost line emitting region, it could be hidden if the central engine is not seen nearly face-on.

4 References

- Boroson T., Green R.: 1992, ApJ 80,109
Burstein D., Heyles C., 1984, ApJS 54, 33
Dumont A.-M., Collin-Souffrin S.: 1990, A& A 229, 313
Ferland G.J., Netzer H., 1983, ApJ 264, 105
Ferland G.J., Korista K.T., Peterson B.M.: 1990, ApJ 363, L121
Ferland G.J., Peterson B.M., Horne K., Welsh W.F., Nahar S.N., 1992, ApJ 387, 95
Gaskell, M. 1982, ApJ 263, 79

Grandi S., 1983, ApJ 268, 591

Hummer D.G., Storey P., 1987, MNRAS

Lightman A.P., White T.R., 1988, ApJ 335, 57

MacAlpine G.M., 1981, ApJ 251, 465

MacAlpine G.M., Davidson K., Gull T.R., Wu C.C., 1985, ApJ 294, 147

Marziani P., Calvani M., Sulentic J.W., 1992, ApJ 393, 658

Morris S.L., Ward M.J., 1989, ApJ 340, 723

Netzer H., Elitzur Ferland G.J., 1983, ApJ 299, 752

Osterbrock D.E., 1977, ApJ 215, 733

Osterbrock, 1988, Astrophysics of Gaseous Nebulae and Active Galactic Nuclei (Mill Valley: Univ. Science Books)

Osterbrock D.E., Shaw R.A., 1988, ApJ 327, 89

Osterbrock D.E., Shuder J. M., 1982, ApJS 49, 149

Netzer H., 1990, in Active Galactic Nuclei, p. 57 (Berlin: Springer).

Peterson B.M., Ferland G.K., 1986, Nature 324, 345

Rees M.J., Netzer H., Ferland G.K., 1989 ApJ 347, 640

Shuder J.M., 1982, ApJ 259, 48

Sulentic J.W. 1989, ApJ 343, 54

Sulentic J.W., Calvani M., Marziani, P., Zheng W., 1990, ApJ 355, L15

Yaqoob T., Warwick R.S., Pounds K.A., 1989 MNRAS 236, 153

Wamsteker W., et al., 1990, ApJ 354, 446

5 Figure Caption

Fig.1: Spectrum of PG 1138+222. Horizontal scale is wavelength in Å; vertical scale is flux in units of $10^{-15} \text{ ergs s}^{-1} \text{ cm}^{-2} \text{ Å}^{-1}$.

Fig.2: Fits to the broad components and to the underlying continuum of the strongest lines: (a) H γ ; (b) HeII λ 4686; (c) H β . Scale units are as for Fig. 1.

Fig.3: Profile ratio between $\text{HeII}\lambda 4686_{BC}$ and $\text{H}\beta_{BC}$. Horizontal scale is radial velocity (binned to 225 km s^{-1}). The origin has been set at the average radial velocity of the narrow components of the two lines.

6 Acknowledgements

P.M. acknowledges the financial support provided by Italian Consiglio Nazionale delle Ricerche for a short post-doctoral period at the University of Alabama. Extragalactic studies at the University of Alabama are supported by NSF/State of Alabama EPSCoR grant RII-8996152.